



The Use of Analgesics in Rodents and Rabbits

Deborah M. Mook, DVM


(Last updated June, 2009)

Rationale.

The Animal Welfare Act (AWA) defines a painful procedure as any procedure that would reasonably be expected to cause more than slight or momentary pain and/or distress in a human being to which that procedure is applied. Emory University's Institutional Animal Care and Use Committee adopts this definition for all species and expects analgesics to be used to alleviate pain unless there is scientific justification to do otherwise.

In addition, well-established studies have shown repeatedly that effective analgesia enhances locomotion, increases appetite, and reduces the time of postoperative recovery for humans, rodents, rabbits, which are particularly sensitive to pain and inflammation, and other species. The figure below, derived from Liles and Flecknell (1999) illustrates the value of post-operative analgesics in improving recovery of rodents in the first 24 hours following laparotomy. Weight loss and decreases in appetite and water consumption were less in rats given an analgesic agent (buprenorphine) than those given saline. As a general rule, for maximal effectiveness, analgesics should be first administered before the animal is fully recovered from anesthesia and should be continued for the next 48-72 hours. However, as little as a single post-operative dose, in many circumstances, is sufficient to dramatically and positively influence recovery.

Effect of Post-op Buprenorphine on Rats



Mechanism of Action. Analgesics can act through several mechanisms. Centrally-acting agents, such as morphine, meperidine, codeine, nalbuphine and buprenorphine, interact with the endogenous opioid systems in the CNS. Substances that act locally at the nociceptor, such as local anesthetics, antihistamines, and alpha-2 adrenoreceptor agonists (i.e. xylazine, clonidine, metomidate), block nerve impulses. Nonsteroidal anti-inflammatory drugs (NSAID), such as aspirin, flunixin meglumine and acetaminophen, inhibit the production of the chemical mediators that activate peripheral nociceptors.

Nonsteroidal Anti-Inflammatory Drugs. NSAIDs are sufficiently potent to treat musculoskeletal, incisional, and acute, mild visceral pain. They inhibit the production of metabolic products of arachidonic acid (prostaglandins, prostacyclin) which are potent chemical mediators of inflammation that activate peripheral nociceptors. Because of different sites of action, these agents are synergistic with opioids. However, combinations of NSAIDs should not be used together as they may cause gastritis/gastric ulceration, or renal failure. Platelet inhibition is also a risk especially with agents with cyclooxygenase I activity. Carprofen (Zenecarp) has low cyclooxygenase 1 inhibitory effects and a low potential for renal or gastric toxicity (safer than flunixin). It is very effective when combined with opioids or buprenorphine and has a 24-48 hour duration of efficacy in most species.

Opioids. For the control of acute or chronic visceral pain, opioids are the most powerful and effective analgesics. Bradycardia, respiratory depression, excessive sedation, nausea, ileus and pica may be side effects. Opioid agonist-antagonists, such as buprenorphine, have minimal side effects. Opioids and opioid agonist-antagonists may increase CSF pressure and should be used with caution in craniotomy cases. In addition, the use of most of these agents is strictly controlled by the federal government requiring specialized licensure and the need to record every drop used. The use of traditional, parenterally-administered opioid analgesics such as morphine, meperidine and pentazocine are impractical in rabbits and rodents because of their high metabolic rates which necessitates intensive dosing schedules to maintain therapeutic blood concentrations. Liposome-encapsulated opioid preparations have been reported in rats, along with recipes for generating them, a formulation that may provide analgesia for as long as 7 days with a single injection (Smith et al., 2003). The administration of NSAIDS concomitantly with opioids may allow the effective use of lower doses of opioids with fewer side effects and adequate pain management.

Oral Administration of Analgesic Agents. Voluntary oral consumption of drugs may be impeded by palatability, xenophobia, pain/discomfort, and recovery from anesthesia and generally is ineffective for immediate post-operative use. In addition, consumption of agents may be influenced by diurnal behavioral rhythms (nocturnal consumption is likely to be greater than that during daylight). Oral doses of these agents may be substantially higher than parenteral doses due to first pass effects through the liver (i.e., oral doses of buprenorphine and butorphanol are 10 times that of injected doses). Many orally administered agents are only adequate for controlling low grade pain and some agents, such as acetaminophen, have little oral potency in rodents. An innovative way of providing oral analgesics to rodents is via flavored gelatin. Grape, currant or cherry-

flavored gelatins containing drugs such as ketoprofen (0.4 mg/ml of gelatin) can be administered to rats and mice. Feed 2 ml of gelatin per 200 gram rat once daily. Feeding the gelatin without drugs for a few days conditions the animals to readily accept the drug-impregnated gelatin later. For some drugs, i.e. buprenorphine, the necessary quantity of drug required makes the gelatin unpalatable. In cases where medication is to be delivered in the drinking water, administration for a few days prior to surgery may address the neophobic response. Ultimately, gavage is the only route by which administration of appropriate doses of oral medications can be ensured.

Common Analgesic Doses (in mg/kg) for Rodents and Rabbits

Agent	Guinea Pig	Mouse	Rat	Rabbit
Acetaminophen	----	----	----	1 ml elixir in 100ml drinking water
Aspirin	87 PO BID	100-120 PO BID	100-120 PO BID	100 PO BID
Buprenorphine	0.05 SC q 8-12 hr	0.05-0.1 SC q 8-12 hr	0.01-0.1 SC q 6-12 hr 5-10 PO q 12 hr	0.01-0.05 SC q 8-12 hr
Butorphanol	----	1-5 SC q 4-6 hr	2 SC q 4-6 hr	0.1-0.5 SC q 4-6 hr
Carprofen (Rimadyl)	----	----	5 SC BID	1.5 PO BID
Flunixin meglumine (Banamine)	----	2.5 SC BID	2.5 SC BID	1.1 SC SID-BID
Ibuprofen	----	40/day in drinking water	15/day in drinking water	----
Ketoprofen	----	----	2 PO SID	3 IM BID

Ketorolac	----	5.0-10.0 PO SID	3-5 PO SID-BID 1 IM SID- BID	----
Meloxicam		1 mg/kg SC or PO q 24 hr*	1 mg/kg SC or PO q 24 hr	
Meperidine	10-20 SC q 2-3 hr	10-20 SC q 2-3 hr	5-10 SC q 2-3 hr	5-10 SC q 2-3 hr; 0.2 mg/ml in drinking water
Morphine	2-5 SC q 2-4 hr	2.5 SC q 2-4 hr	2.5 SC q 2-4 hr	2-5 SC q 2-4 hr
Morphine (liposome encapsulated)			2.8 SC once	
Nalbuphine	1-2 q 4-8 hr	4-8 SC q 4-8 hr	1-2 SC q 4-8 hr	1-2 SC q 4-8 hr
Oxymorphone	----	0.1 SC q 4 hr	0.3 SC q 4 hr	----
Oxymorphone (liposome encapsulated)			1.2 SC once	
Pentazocine	----	10 SC q 3-4 hr	10 SC q 4 hr	5-10 SC q 2-4 hr
Xylazine	----	-----	5-12 SC q 2 hr	----

In mice, it is recommended to dilute injectable meloxicam to a concentration of 0.5 mg/ml.

Selected References:

Chellman GJ, et al. 1994. Comparison of ketorolac tromethamine with other injectable nonsteroidal anti-inflammatory drugs for pain-on-injection and muscle damage in the rat. **Human Exp Toxicol** 13: 111-7.

- Flecknell P.A. 1996. Anesthesia and analgesia for rodents and rabbits. In: **Handbook of Rodent and Rabbit Medicine**, Laber-Laird K, Swindle MM and Flecknell PA, eds., Pergamon Press, Butterworth-Heinemann, Newton, MA, pp. 219-37.
- Flecknell, P.A. 1996. **Laboratory Animal Anaesthesia: An Introduction for Research Workers and Technicians**. 2nd ed., Academic Press, London, U.K.
- Flecknell PA, Orr HE, **Roughan** JV, Stewart R. 1999. **Comparison of the effects of oral or subcutaneous carprofen or ketoprofen in rats undergoing laparotomy**. **Veterinary Record**. 144(3):65-7.
- Granados-Soto V, et al. 1995. Relationship between pharmacokinetics and the analgesic effect of ketorolac in the rat. **J Pharmacol Exp Therap** 272: 352-6.
- Hayes JH, Flecknell PA. 1999. A comparison of pre- and post-surgical administration of bupivacaine or buprenorphine following laparotomy in the rat. **Lab Animals** 33: 16-23.
- Kelley MC, et al. Ketorolac prevents postoperative intestinal ileus in rats. **Amer J Surg** 165: 107-11.
- Liles JH, Flecknell PA. 1992a. The effects of buprenorphine, nalbuphine and butorphanol alone or following halothane anaesthesia in food and water consumption and locomotor movement in rats. **Lab Animals** 26: 180-9.
- Liles JH, Flecknell PA. 1992b. The use of non-steroidal anti-inflammatory drugs for the relief of pain in laboratory rodents and rabbits. **Lab Animals** 26: 242-55.
- Liles JH, Flecknell PA. 1993a. The influence of buprenorphine or bupivacaine on the post-operative effects of laparotomy and bile-duct ligation in rats. **Lab Animals** 27: 374-80.
- Liles JH, Flecknell PA. 1993b. The effects of surgical stimulus on the rat and the influence of analgesic treatment. **Br Vet J** 149: 515-25.
- Liles JH, Flecknell PA. 1994. A comparison of the effects of buprenorphine, carprofen and flunixin following laparotomy in rats. **J Vet Pharmacol Therap** 17: 284-90.
- Liles JH, Flecknell PA, et al. 1998. Influence of oral buprenorphine, oral naltrexone or morphine on the effects of laparotomy in the rat. **Lab Animals** 32: 149-61.
- Martin, L.B et al. 2001. Analgesic efficacy of orally administered buprenorphine in rats. **Comparative Medicine** 51(1): 43-48.
- Martin, T.J. et al. 2004. Effects of laparotomy on spontaneous exploratory activity and conditioned operant responding in the rat. **Anesthesiology** 101(1): 191-203.
- Roughan, J.V., Flecknell, P.A. & Davies, B.R. Behavioral assessment of the effects of tumor growth in rats and the influence of the analgesics carprofen and meloxicam. **Lab. Anim.** 2004 38(3): 286-96.
- Roughan, J.V. & Flecknell, P.A. 2003. Evaluation of a short duration behavior-based post-operative pain scoring system in rats. **European Journal of Pain** 7(5): 397-406.
- Smith, L.J. et al. 2003. A single dose of liposome-encapsulated oxymorphone or morphine provides long-term analgesia in an animal model of neuropathic pain. **Comparative Medicine** 53 (3): 280-287.

Stewart, L. et al. 2003. Evaluation of postoperative analgesia in a rat model of incisional pain. Contemporary Topics 42(1): 28-34.

Wootton R, Cross G, et al. 1988. An analgesiometry system for use in rabbits with some preliminary data on the effects of buprenorphine and lofentanil. **Lab Animals** 22: 217-22.

[Return to Main](#)

Copyright © Emory University, 1997. All Right reserved.
Send Comments to postmaster@dar.emory.edu